

## Grapiprant

### - Drug Information

#### **Brand Names:**

Galliprant®

#### **Pharmacology:**

A non-cyclooxygenase inhibiting, non-steroidal anti-inflammatory drug

#### **Indications:**

Labeled for the control of pain and inflammation associated with osteoarthritis in dogs.

#### **Contraindications:**

##### **- General**

For oral use in dogs only.

Dogs less than 8 pounds. (3.6 kgs) cannot be accurately dosed.

Should not be used in dogs that have a hypersensitivity to grapiprant.

The safe use has not been evaluated in dogs younger than 9 months of age, dogs used for breeding, or in pregnant or lactating dogs.

Grapiprant is a methylbenzene sulfonamide. It is not known whether dogs with a history of hypersensitivity to sulfonamide drugs will exhibit hypersensitivity to grapiprant.

#### **Interactions:**

##### **- General Interactions**

No identified General Interactions at time of publishing.

##### **- Category Interactions**

- Corticosteroid

Contraindicated with concurrent use of steroids because it may increase the risk of side effects.

- NSAID-Non Steroidal Antiinflammatory Drug

Contraindicated with concurrent use of other NSAIDs because it may increase the risk of side effects.

- Protein bound drug

Concomitant use of protein-bound drugs has not been studied.

##### **- Drug Interactions**

#### **Adverse Effects:**

##### **- General**

Adverse reactions may include vomiting, diarrhea, decreased appetite, mucoid, watery or bloody stools, and decreases in serum albumin and total protein.

#### **Dosages:**

##### **- Small Animal**

- Dog

2 mg/kg by mouth once daily

Tablets are scored and dosage should be calculated in half tablet increments.

#### **Special Notes:**

None